

Probing the gravitational redshift effect from the relativistic jets of compact AGN

Tigran G. Arshakian^{1,*}

Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany

Abstract. I explore a possibility to measure the gravitational redshift (GR) effect in the gravitational field of massive central nuclei residing in active galaxies (AGN). The activity of central nuclei is associated with the bipolar jet ejection of relativistic plasma which produces strong radio emission. I consider the behavior of the flux density variations of the jet plasma as a result of GR effect from a Schwarzschild black hole, and I discuss possibilities to detect the GR effect from the relativistic jets of compact AGN with present and future radio facilities.

1 Introduction

Development of new astronomical facilities provides high resolution and sensitivity thus allowing one to reach the scales where the General Relativity effects can be tested. There is already compelling evidence that active galaxies host super massive black holes (SMBH) reaching up to $\sim 10^{10}$ solar masses. The central active nuclei are associated with the relativistic jets which originate near the central nuclei and move out with relativistic speeds exceeding $0.9c$. The jet plasma radiates strong synchrotron radio emission which is detectable on parsec-scales by very large baseline interferometry (VLBI). Here, we consider the flux-density variation of the jet plasma due to the GR effect, and the possibility of its detection by present and future radio-astronomical facilities.

2 Flux variation of the relativistic jet

Suppose that the jet plasma component radiating a power law spectrum $F_e(\nu) \sim \nu^{-\alpha}$ (α is the spectral index) moves in the gravitational field of SMBH with a speed β in a direction with an angle θ to the line of sight of the observer. Then the observed flux at a time t depends on the Doppler factor $\delta_t(\beta; \theta)$ of the component and the distance of the component from the SMBH at t , R_t ,

$$F_t(\nu) = k_t^{3+\alpha} F_e(\nu); \quad k_t = \frac{\delta_t}{1+z} \left(1 - \frac{R_s}{R_t}\right)^{1/2}; \quad (1)$$

where R_s and z are the Schwarzschild radius of the SMBH and its redshift. The detected flux density may vary as a result of deceleration or acceleration

* On leave from Byurakan Astrophysical Observatory, Byurakan 378433, Armenia

of the jet component, bending of the jet and intrinsic or extrinsic variability of radio emission. To investigate the pure effect of the GR on the flux density variation, we assume, that the $F_e(\nu)$ and δ_t remain unchangeable in time. At a fixed frequency ν_f , the detected flux density $F(\nu_f)$ of the jet component grows with R_t (Fig. 1) reaching up to 90 % of its intrinsic flux density at $\sim 20R_s$, and then gradually approaches to ~ 100 % at $\sim 100R_s$.

3 Prospects of detecting the GR

Seven millimeter VLBI¹ (43 GHz) can achieve an angular resolution of $\sim 70 \mu\text{as}$ corresponding to a spatial resolution of $0.0058 \text{ pc} \simeq 19R_s$ for M87². Full coverage global mm-VLBI imaging with more sensitive radio telescopes will allow even more detailed and better quality images¹. Present high-resolution multi-epoch mm-VLBI imaging of M87 is capable of detecting ~ 10 % flux density variations (Fig. 1) from the relativistic jet plasma at $\sim (15 \text{ to } 100)R_s$ due only to GR effect.

At lower frequency, a global VLBI array enhanced by the SKA will allow one to study the nearest weak AGN which previously were unreachable by VLBI due to their faintness. The next generation space-VLBI mission VSOP-2³ will achieve the highest resolution of $38 \mu\text{as}$ at 43 GHz, which is twice as high ($\sim 10R_s$ for M87) as the resolution of a present mm-VLBI. The resolution of a global sub-mm-VLBI¹ at 230 GHz will be $< 30 \mu\text{as}$ which is comparable with the resolution of future space-VLBI missions. The global sub-mm-VLBI combined with ALMA can achieve the unprecedented angular resolution of $\sim 10 \mu\text{as}$ ($\sim 3R_s$ for M87!) allowing the gravitational effects to be tested in the vicinity of SMBH thus providing additional test for the theory of general relativity and a new method for measuring the masses of SMBH.

TGA is grateful to the AvH Foundation for a Humboldt Fellowship, and to Drs. T. Krichbaum, A. Lobanov and A. Polatidis for useful discussions.

¹ Krichbaum et al., poster #43 in this proceedings (see also: <http://www.mpifr-bonn.mpg.de/staff/tkrichbaum/3c274.html>)

² M87 (3C 274) has a radio bright core-jet structure, $z = 0.0044$ ($D \sim 17.1 \text{ Mpc}$, for a flat cosmology with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$), $M_{\text{BH}} \sim 3 \times 10^9 M_\odot$ ($R_s = 0.0003 \text{ pc}$)

³ Hirabayashi et al., poster #4 in this proceedings




Fig. 1. The observed flux density normalized to the intrinsic flux density of the jet component versus its distance (in R_s units) from the SMBH. Amplification of the flux density due to GR effect is shown for a flat $\alpha = 0$ (solid line) and steep $\alpha = 1$ (dashed line) spectra.